



Mapping of high-elevation alpine grassland communities based on hyperspectral UAV measurements

Levente Papp^{1,2}, Abraham Mejia-Aguilar², Ruth Sonnenschein², Rita Tonin³, Michael Loebmann³, Clemens Geitner⁴, Martin Rutzinger⁴, Andreas Mayr⁴ and Stefan Lang¹

¹ Department of Geoinformatics - Z_GIS, University of Salzburg, Salzburg, Austria

² EURAC Research, Bolzano, Italy



³ Faculty of Science and Technology, Free University of Bolzano, Bolzano, Italy

⁴ Department of Geography, University of Innsbruck, Innsbruck, Austria



Study area

- Funes valley, in Puez-Geisler National Park, South Tyrol, Italy
- 2190-2300 m a.s.l.
→ precipitous area
- Endangered by shallow erosion




Funes valley study area near-infrared composite (896 nm)

- Area: 5 ha
- Field trips:
 - 23/08/2019
 - Botanical surveys
 - 04/09/2019
 - UAV flights
 - Field measurements




Funes valley study area RGB composite



Funes valley study area digital elevation model

Measurements



UAV flights



RGB image



Hyperspectral image: Rikola camera:
40 bands: 506-896 nm
5 cm spatial accuracy

Field measurements (Quadrats) – 50x50 cm:



3 measurements with Spectroradiometer (Spectra Vista HR-1024i)
from 1 meter high

Range: 340-2500 nm

(+ Photos with spectroradiometer, for positioning)



High quality RGB images





Hyperspectral images of quadrats




Sampled vegetation

+ GNSS RTK GMPS measurements of the corner of quadrats



Measurements


Hyperspectral UAV measurements



Excluded bands:
(12)

Band 26 – 756,129 nm
Band 27 – 766,380 nm
Band 28 – 776,245 nm
Band 29 – 786,129 nm
Band 30 – 796,154 nm
Band 31 – 806,339 nm
Band 33 – 825,500 nm
Band 34 – 835,804 nm
Band 35 – 846,697 nm
Band 36 – 855,982 nm
Band 37 – 866,602 nm
Band 38 – 877,456 nm


28 bands remaining for the classification



Workflow


Field

Lab



Defined classes

Funes valley study area – defined vegetation classes








Legend



Quadrats

Defined vegetation classes

-  Class 01 – *Anthoxanthum alpinum* community
-  Class 02 – *Avenula pubescens* & *Festuca norica* community
-  Class 03 – *Avenula pubescens* & *Triestum flavescens* community
-  Class 04 – Short grass
-  Class 05 – *Geranium sylvaticum* community

25 0 25 50 75 100 m
Created by Levente Papp, – EURAC Center for Sensing Solutions, 2020 – ETRS89/UTM zone N32


Defined classes

- Based on the botanical survey on the first field trip

Class 01 – Anthoxanthum alpinum community



A very bright green and dense grass, in the surveyed plots, it is really homogenous. The areas were clearly defined on the ground and on the orthomosaic too.





| | |
|------------------------------------|-------|
| <u>Anthoxanthum alpinum</u> | Grass |
| <u>Phleum rhaeticum</u> | Grass |
| <u>Avenula pubescens</u> | Grass |
| Festuca nigrescens | Grass |
| Geraneum sylvaticum | Forb |
| Knautia longifolia | Forb |
| Trollius europaeus | Forb |
| Achillea millefolium | Forb |
| Pulsatilla alpina | Forb |
| Carduus defloratus | Forb |
| Pimpinella major | Forb |
| Silene vulgaris | Forb |
| Carex sempervirens | Grass |
| Poa glauca | Grass |
| Hypericum maculatum | Forb |




Defined classes

Class 01 – Anthoxanthum alpinum community



66% - Anthoxanthum alpinum


| | |
|--------------------------|-------|
| <u>Phleum rhaeticum</u> | Grass |
| <u>Avenula pubescens</u> | Grass |
| Festuca nigrescens | Grass |
| Geraneum sylvaticum | Forb |
| Knautia longifolia | Forb |
| Trollius europaeus | Forb |
| Achillea millefolium | Forb |
| Pulsatilla alpina | Forb |
| Carduus defloratus | Forb |
| Pimpinella major | Forb |
| Silene vulgaris | Forb |
| Carex sempervirens | Grass |
| Poa glauca | Grass |
| Hypericum maculatum | Forb |



Spectral signals of the quadrats measured with the spectroradiometer

Class 02 – *Avenula pubescens* and *Festuca norica* community

One of the most common species on the study site. This yellowish-green plants also occurs in all of the other classes, but are only in this area the dominant species.





| | |
|---------------------------------|--------|
| <u><i>Avenula pubescens</i></u> | Grass |
| <u><i>Festuca norica</i></u> | Grass |
| <u><i>Avenula praeusta</i></u> | Grass |
| <i>Sesleria varia</i> | Grass |
| <i>Scabiosa lucida</i> | Forb |
| <i>Trollius europaeus</i> | Forb |
| <i>Arnica montana</i> | Forb |
| <i>Trifolium pratense</i> | Legume |
| <i>Carduus defloratus</i> | Forb |
| <i>Leontodon hispidus</i> | Forb |
| <i>Horminum</i> | |
| <i>pyrenaicum</i> | Forb |
| <i>Achillea millefolium</i> | Forb |
| <i>Pulmonaria australis</i> | Forb |
| <i>Briza media</i> | Grass |



Defined classes


Class 02 – *Avenula pubescens* and *Festuca norica* community



No collected vegetation samples


70% - *Avenula pubescens* & *Festuca norica*

| | |
|-----------------------------|--------|
| <i>Avenula praeusta</i> | Grass |
| <i>Sesleria varia</i> | Grass |
| <i>Scabiosa lucida</i> | Forb |
| <i>Trollius europaeus</i> | Forb |
| <i>Arnica montana</i> | Forb |
| <i>Trifolium pratense</i> | Legume |
| <i>Carduus defloratus</i> | Forb |
| <i>Leontodon hispidus</i> | Forb |
| <i>Horminum pyrenaicum</i> | Forb |
| <i>Achillea millefolium</i> | Forb |
| <i>Pulmonaria australis</i> | Forb |
| <i>Briza media</i> | Grass |




Spectral signals of the quadrats measured with the spectroradiometer

Class 03 – Avenula pubescens and Trisetum flavescens community







This community mostly occurs in the humid valley of the area, it is slightly darker than its surroundings.

| | |
|----------------------------|--------|
| <u>Trisetum flavescens</u> | Grass |
| <u>Avenula pubescens</u> | Grass |
| <u>Phleum rhaeticum</u> | Grass |
| Achillea millefolium | Forb |
| Rumex alpestris | Forb |
| Geraneum sylvaticum | Forb |
| Knautia longifolia | Forb |
| Vicia cracca | Legume |
| Hypericum maculatum | Forb |
| Alchemilla hirsuta group | Forb |
| Trollius europaeus | Forb |
| Leontodon hispidus | Forb |
| Silene vulgaris | Forb |
| Pimpinella major | Forb |



Defined classes

Class 03 – *Avenula pubescens* and *Trisetum flavescens* community



Spectral signals of the quadrats
measured with the spectroradiometer

56% - *Avenula pubescens* & *Trisetum flavescens*

| | |
|-------------------------|--------|
| <u>Phleum rhaeticum</u> | Grass |
| Achillea millefolium | Forb |
| Rumex alpestris | Forb |
| Geraneum sylvaticum | Forb |
| Knautia longifolia | Forb |
| Vicia cracca | Legume |
| Hypericum maculatum | Forb |
| Alchemilla hirsuta | |
| group | Forb |
| Trollius europaeus | Forb |
| Leontodon hispidus | Forb |
| Silene vulgaris | Forb |
| Pimpinella major | Forb |

Class 04 – Short grass

Originally, we defined this as an independent vegetation class, but after a more detailed survey, we found these areas are homogenous because of anthropogenic influences. – No recorded quadrats because of this.



| | |
|----------------------------|--------|
| <u>Trisetum flavescens</u> | Grass |
| <u>Carex sempervirens</u> | Grass |
| <u>Sesleria varia</u> | Grass |
| Carex caryophylla | Grass |
| Luzula alpina | Grass |
| Phyteuma orbiculare | Forb |
| Carduus defloratus | Forb |
| Hieracium morisianum | Forb |
| Scabiosa sp | Forb |
| Gentiana anisodonta | Forb |
| Aster bellidiastrum | Forb |
| Horminum pyrenaicum | Forb |
| Anthyllis vulneraria | Legume |
| Pedicularis elongata | Forb |
| Festuca norica | Grass |
| Polygonum viviparum | Forb |



Class 05 – Geranium sylvaticum community



Rusty coloured in this season (September)
– characteristic field pattern, occurs in a variety of locations in the whole study area.

No recorded quadrats

| | |
|----------------------------------|--------|
| <u>Geranium sylvaticum</u> | Forb |
| <u>Avenula pubescens</u> | Grass |
| <u>Festuca norica</u> | Grass |
| <u>Alchemilla group hirsutae</u> | Forb |
| Trifolium pratense | Legume |
| Trollius europaeus | Forb |
| Sesleria varia | Forb |
| Phleum rhaeticum | Grass |
| Silene vulgaris | Forb |
| Leontodon hispidus | Forb |
| Rumex alpestris | Forb |
| Achillea millefolium | Forb |
| Ranunculus nemorosus | Forb |
| Pulsatilla alpina | Forb |




Other defined classes

Modified and non-vegetation classes:

- Sparse/replanted areas
In some areas they replanted the vegetation, spectrally close to the bare soil because it is not dense enough to cover it
- Covered areas
Same, but covered with textile and cut-of grass (Class 4)
- Wooden barriers
Near the replanted areas to stop the erosion
- Drooped (or trampled) grass
The areas where traces from a field survey two weeks earlier that were still visible
- Shadow
- Bare soil
- Bare rock

Σ **12 classes** (5 grassland communities + 7 „others”)




Classification

For training:


- Based on the botanical survey in the first field trip
- Smaller subsets of it based on the orthomosaic
- Delineated homogenous areas with high precision (GNSS RTK GPS)
- Non-vegetation classes based on orthomosaic – 2/3 for training

For validation:



- 50x50 cm quadrats - knowing the exact content of them
- Areas: approximately 350-400 pixels
- Corner points measured with high precision (GNSS RTK GPS)
- Non-vegetation classes based on orthomosaic – 1/3 for validation
- Validation with Confusion matrix



Challenges




Spectroradiometer




PCA – Principal Component Analysis

- Based on Eigenvalues we reduced the variables
- 28 bands reduced to **6 bands** which contains the majority of the information



Support Vector Machine classification

- A supervised learning method
- Try to find the proper hyperplanes in the spectral space between the classes
- Suitable for smaller datasets (e.g. not like ANN)
- We set the proper parameters based on one classification rule image (the probability of a given pixel belongs to that class)
- Used software: ENVI Classic 5.2
















Source: G.Mountrakis et.al. (2010)

Result map

Support Vector Machine Classification of Funes Valley study area

Legend




-  *Anthoxanthum alpinum* community
-  *Avenula pubescens* & *Festuca norica* c.
-  *Avenula pubescens* & *Trisetum flavescens* c.
-  Short grass
-  *Geranium sylvaticum* community
-  Sparse vegetation / re-planted areas
-  Bare soil
-  Wooden barriers
-  Covered areas
-  Drooped grass
-  Shadow
-  Bare rock


0 25 50 m

Created by Levente Papp, Abraham Mejia-Aguilar, Ruth Sonnenschein, Rita Tonin
– EURAC, UNIBZ, 2020
ETRS89/UTM zone N32


Examples




Humid Valley with Class 03 – *Avenula pubescens* and *Trisetum flavescens* community and drooped grass



A shallow erosion area with sparsen vegetation surrounded by Class 05: *Geranium sylvaticum* and a combination of the two *Avenula pubescens* classes (Class 02 and Class 03)



Shallow erosion areas (covered and replanted ones with the wooden barriers) – surrounded by Class 04: short grass and Class 05: *Geranium sylvaticum*



Results

| Classes: | 01 | 02 | 03 | 04 | 05 |
|-----------------------------|-----|-----|-----|----|----|
| Achillea millefolium | | | | | |
| Alchemilla hirsuta group | | | | | |
| Anthoxanthum alpinum | 66% | <5% | | | |
| Anthyllis vulneraria | | | | | |
| Arnica montana | | | | | |
| Aster bellidiastrum | | | | | |
| Avenula praeusta | | | | | |
| Avenula pubescens | | 50% | 20% | | |
| Briza media | | <5% | | | |
| Carduus defloratus | | | | | |
| Carex caryophyllea | | | | | |
| Carex sempervirens | | | | | |
| Chaerophyllum hirsutum | | | | | |
| Festuca nigrescens | 10% | | | | |
| Festuca norica | | 20% | | | |
| Gentiana anisodonta | | | | | |
| Geraneum sylvaticum | 7% | | 7% | | |
| Hieracium morisianum | | | | | |
| Horminum pyrenaicum | | | | | |
| Hypericum maculatum | | | | | |

| | | | | | |
|----------------------------|--|-----|-----|--|--|
| Knautia longifolia | | | | | |
| Leontodon hispidus | | | | | |
| Luzula alpina | | | | | |
| Pedicularis elongata | | | | | |
| Phleum rhaeticum | | | | | |
| Phyteuma orbiculare | | | | | |
| Pimpinella major | | | 7% | | |
| Poa glauca | | | | | |
| Polygonum viviparum | | | | | |
| Pulmonaria australis | | | | | |
| Pulsatilla alpina | | | | | |
| Ranunculus nemorosus | | | | | |
| Rumex alpestris | | | | | |
| Scabiosa lucida | | | | | |
| Scabiosa sp | | | | | |
| Sesleria varia | | | | | |
| Silene vulgaris | | | | | |
| Trifolium pratense | | | | | |
| Trisetum flavescens | | | 36% | | |
| Trollius europaeus | | <5% | | | |
| Vicia cracca | | | 20% | | |

- One of the dominant species
- Occurs in the area but in minor
- Does not occur in the area
- Percentage coverage inside the community (only if it is covered by a quadrat)

Validation

- The classified map – confusion matrix & the field expert


Overall Accuracy = (2001/2648) 75.5665%
Kappa Coefficient = 0.6785

- Overrepresented/false positive results?

- Small misclassification in case of class 2/3
- Reasons: Species overlapping **Avenula pubescens** but they are two different communities
- For the community of class 3: *Avenula pubescens* & *Trisetum flavescens* the time of the field measurements (September) quite late. The plants are starting yellowing during September and become similar to class 2.

- Geranium sylvaticum* (Class 05) – beside of *Avenula pubescens* – Is the most common species, this community surrounding the other classes
- Class 4 (Short grass): Surrounding the shallow erosion plots and in the western cultivated areas - combined with drooped grass, classified as our result because the slope & anthropogenic influence


| Confusion matrix - Ground truth (percent) | | | | | | |
|---|----------|----------|----------|----------|----------|-------|
| | Class 01 | Class 02 | Class 03 | Class 04 | Class 05 | Total |
| Class01 | 61.61 | 0 | 0.78 | 0 | 0 | 7.33 |
| Class02 | 0 | 32.19 | 0 | 0 | 2.88 | 5.29 |
| Class03 | 38.39 | 35.36 | 89.3 | 0 | 0 | 22.47 |
| Class04 | 0 | 1.06 | 7.83 | 80.15 | 3.69 | 30.97 |
| Class05 | 0 | 31.4 | 2.09 | 19.85 | 93.43 | 33.95 |
| Total | 100 | 100 | 100 | 100 | 100 | 100 |




Discussion

- The major communities were delineated and investigated in two different scales
- A database was built up on species-level for the classes/communities
 - The dominant species and their coverage has been defined
- Hyperspectral aerial image was created and classified with high accuracy
- The classification represents the main field conditions and patterns
- Smaller misclassification on a minor area between class 2 and class 3 caused by the late investigation date

The grassland communities were mapped based on high precision ground measurements and hyperspectral remotely sensed datasets





Thank you for your attention!

Mapping of high-elevation alpine grassland communities based on hyperspectral UAV measurements

The research leading to these results has received funding from the Province of Bolzano under the Research and Innovation action, L§14 and from the European Regional Development Fund, Operational Program Investment for growth and jobs ERDF 2014-2020 under Project number ERDF1094, Data Platform and Sensing Technology for Environmental Sensing LAB-DPS4ESLAB.

